

PERITONEAL MALIGNANT MESOTHELIOMA Piemonte, 1990-2017

Peritoneal malignant mesothelioma. Observed cases, standardized incidence rates (per 100,000 per year; standard: European population, EU27) and 95% confidence intervals (CI) by calendar period and sex.

INCIDENCE 1990-2017 (definite, probable and possible diagnoses)						
	MEN			WOMEN		
Period	observed	rates	95% CI	observed	rates	95% CI
1990-1994	34	0.3	0.2-0.5	17	0.1	0.1-0.2
1995-1999	40	0.4	0.3-0.5	42	0.3	0.2-0.4
2000-2004	40	0.4	0.3-0.5	42	0.3	0.2-0.4
2005-2009	47	0.4	0.3-0.5	35	0.3	0.2-0.4
2010-2014	33	0.3	0.2-0.4	26	0.2	0.1-0.3
2015-2017	20	0.3	0.2-0.4	10	0.1	0.0-0.2

INCIDENCE 1990-2017 (definite diagnoses)						
	MEN			WOMEN		
Period	observed	rates	95% CI	observed	rates	95% CI
1990-1994	26	0.3	0.2-0.4	12	0.1	0.0-0.2
1995-1999	33	0.3	0.2-0.4	32	0.3	0.2-0.4
2000-2004	25	0.2	0.1-0.3	23	0.2	0.1-0.3
2005-2009	31	0.3	0.2-0.4	22	0.2	0.1-0.2
2010-2014	27	0.2	0.1-0.3	21	0.2	0.1-0.2
2015-2017	17	0.2	0.1-0.3	9	0.1	0.0-0.2

Peritoneal malignant mesothelioma. Observed cases, standardized incidence rates (per 100,000 per year; standard: European population, EU27) and 95% confidence intervals (CI) by calendar period and local health authority (2013-2017) .

	INCIDENCE 2013-2017 (definite, probable and possible diagnoses)					
local health authority	MEN			WOMEN		
	observed	rates	95% CI	observed	rates	CI 95%
AL	8	0.6	0.2-1.0	2	0.1	0.0-0.3
AT	2	0.4	0.0-0.9	0	0.0	0.0-0.0
BI	0	0.0	0.0-0.0	0	0.0	0.0-0.0
CN1	1	0.1	0.0-0.3	2	0.1	0.0-0.3
CN2	1	0.2	0.0-0.6	0	0.0	0.0-0.0
NO	2	0.2	0.0-0.6	0	0.0	0.0-0.0
TO	5	0.2	0.1-0.4	0	0.0	0.0-0.0
TO3	4	0.3	0.0-0.5	10	0.5	0.2-0.9
TO4	3	0.1	0.0-0.4	2	0.2	0.0-0.4
TO5	2	0.2	0.0-0.6	1	0.1	0.0-0.3
VC	1	0.2	0.0-0.5	1	0.2	0.0-0.5
VCO	2	0.3	0.0-1.1	1	0.2	0.0-0.6

Peritoneal malignant mesothelioma. Observed cases, standardized incidence rates (per 100,000 per year; standard: European population EU27) and 95% confidence intervals (CI) by area and sex. All diagnoses

LOCAL DISTRICT	INCIDENCE 2013-2017 (definite, probable and possible diagnoses)							
	WOMEN			MEN				
	rates	95% CI	observed	rates	95% CI	observed		
Regione Piemonte	0.1	0.1	0.2	19	0.3	0.2	0.4	31
AL/1 (Acqui Terme)	0	0	0	0	0	0	0	0
AL/2 (Alessandria)	0	0	0	0	0.6	0	1.3	2
AL/3 (Casale Monferrato)	0	0	0	0	2.1	0.2	3.9	5
Comune Casale Monferrato	0	0	0	0	3.5	0	7.6	3
AL/4 (Novi Ligure)	0	0	0	0	0	0	0	0
AL/5 (Ovada)	0.9	0	2.6	1	1.2	0	3.5	1
AL/6 (Tortona)	0	0	0	0	0	0	0	0
AL/7 (Valenza)	1.1	0	3.2	1	0	0	0	0
AT/1 (Asti centro)	0	0	0	0	0	0	0	0
AT/2 (Asti nord)	0	0	0	0	0.6	0	1.7	1
AT/3 (Asti sud-Nizza M.to)	0	0	0	0	0.6	0	1.9	1
BI/1(Biella)	0	0	0	0	0	0	0	0
BI/2 (Cossato)	0	0	0	0	0	0	0	0
CN1/1 (Borgo San Dalmazzo)	0	0	0	0	0	0	0	0
CN1/2 (Ceva)	0	0	0	0	0	0	0	0
CN1/3 (Cuneo)	0	0	0	0	0	0	0	0
CN1/4 (Fossano)	0	0	0	0	0	0	0	0
CN1/5 (Mondovì)	0	0	0	0	0	0	0	0
CN1/6 (Saluzzo)	0.8	0	1.8	2	0.5	0	1.4	1
CN2/1 (Alba)	0	0	0	0	0.3	0	1.0	1
CN2/2 (Bra)	0	0	0	0	0	0	0	0
NO/1 (Arona)	0	0	0	0	0	0	0	0
NO/2 (Borgomanero)	0	0	0	0	1.1	0	2.5	2
NO/3 (Galliate-Trecate)	0	0	0	0	0	0	0	0
NO/4 (Novara)	0	0	0	0	0	0	0	0
TO1-10 (Torino)	0	0	0	0	0.2	0	0.4	5
TO3/1 (Collegno)	2.4	0.6	4.2	7	0.7	0	1.8	2
TO3/2 (Giaveno)	0	0	0	0	0	0	0	0
TO3/3 (Orbassano)	0.3	0	0.9	1	0	0	0	0
TO3/4 (Pinerolo)	0	0	0	0	0	0	0	0
TO3/5 (Rivoli)	0.4	0	1.2	1	0	0	0	0
TO3/6 (Susa)	0	0	0	0	0.8	0	1.9	2
TO3/7 (Val Pellice)	1.6	0	4.7	1	0	0	0	0
TO3/8 (Val Chisone e Germanasca)	0	0	0	0	0	0	0	0
TO3/9 (Venaria)	0	0	0	0	0	0	0	0
TO4/1 (Ciriè)	0	0	0	0	0	0	0	0
TO4/2 (Chivasso)	0.4	0	1.3	1	1.2	0	3.0	2
TO4/3 (Settimo Torinese)	0.5	0	1.6	1	0	0	0	0
TO4/4 (San Mauro Torinese)	0	0	0	0	0	0	0	0
TO4/5 (Ivrea)	0	0	0	0	0.3	0	0.8	1
TO4/6 (Cuorgnè)	0	0	0	0	0	0	0	0
TO5/1(Carmagnola)	0	0	0	0	0	0	0	0
TO5/2 (Chieri)	0	0	0	0	0.7	0	1.6	2
TO5/3 (Moncalieri)	0	0	0	0	0	0	0	0
TO5/4 (Nichelino)	0.4	0	1.3	1	0	0	0	0
VC/1(Vercelli-Santhià)	0.3	0	1	1	0	0	0	0
VC/2 (Montano-Borgosesia-Varallo)	0	0	0	0	0.4	0	1.2	1
VCO/1(Domodossola)	0	0	0	0	0	0	0	0
VCO/2 (Omegna)	0	0	0	0	0	0	0	0
VCO/3 (Verbania)	0.5	0	1.5	1	1.2	0	2.8	2

Peritoneal malignant mesothelioma. Observed cases, standardized incidence rates (per 100,000 per year; standard: European population EU27) and 95% confidence intervals (CI) by area and sex. Definite diagnoses

INCIDENCE 2013-2017 (definite diagnoses)							
LOCAL DISTRICT	WOMEN			MEN			
	rates	95% CI	observed	rates	95% CI	observed	
Regione Piemonte	0.1	0.1	0.2	14	0.2	0.1	0.3
AL/1 (Acqui Terme)	0	0	0	0	0	0	0
AL/2 (Alessandria)	0	0	0	0	0.6	0	1.3
AL/3 (Casale Monferrato)	0	0	0	0	2.1	0.2	3.9
Comune Casale Monferrato	0	0	0	0	3.5	0	7.6
AL/4 (Novi Ligure)	0	0	0	0	0	0	0
AL/5 (Ovada)	0.9	0	2.6	1	1.2	0	3.5
AL/6 (Tortona)	0	0	0	0	0	0	0
AL/7 (Valenza)	0	0	0	0	0	0	0
AT/1 (Asti centro)	0	0	0	0	0	0	0
AT/2 (Asti nord)	0	0	0	0	0.6	0	1.7
AT/3 (Asti sud-Nizza M.to)	0	0	0	0	0.6	0	1.9
BI/1(Biella)	0	0	0	0	0	0	0
BI/2 (Cossato)	0	0	0	0	0	0	0
CN1/1 (Borgo San Dalmazzo)	0	0	0	0	0	0	0
CN1/2 (Ceva)	0	0	0	0	0	0	0
CN1/3 (Cuneo)	0	0	0	0	0	0	0
CN1/4 (Fossano)	0	0	0	0	0	0	0
CN1/5 (Mondovì)	0	0	0	0	0	0	0
CN1/6 (Saluzzo)	0.5	0	1.3	1	0	0	0
CN2/1 (Alba)	0	0	0	0	0.3	0	1.0
CN2/2 (Bra)	0	0	0	0	0	0	0
NO/1 (Arona)	0	0	0	0	0	0	0
NO/2 (Borgomanero)	0	0	0	0	1.1	0	2.5
NO/3 (Galliate-Trecate)	0	0	0	0	0	0	0
NO/4 (Novara)	0	0	0	0	0	0	0
TO1-10 (Torino)	0	0	0	0	0.2	0	0.4
TO3/1 (Collegno)	2.1	0.4	3.7	6	0.7	0	1.8
TO3/2 (Giaveno)	0	0	0	0	0	0	0
TO3/3 (Orbassano)	0.3	0	0.9	1	0	0	0
TO3/4 (Pinerolo)	0	0	0	0	0	0	0
TO3/5 (Rivoli)	0	0	0	0	0	0	0
TO3/6 (Susa)	0	0	0	0	0.8	0	1.9
TO3/7 (Val Pellice)	0	0	0	0	0	0	0
TO3/8 (Val Chisone e Germanasca)	0	0	0	0	0	0	0
TO3/9 (Venaria)	0	0	0	0	0	0	0
TO4/1 (Ciriè)	0	0	0	0	0	0	0
TO4/2 (Chivasso)	0.4	0	1.3	1	1.2	0	3.0
TO4/3 (Settimo Torinese)	0.5	0	1.6	1	0	0	0
TO4/4 (San Mauro Torinese)	0	0	0	0	0	0	0
TO4/5 (Ivrea)	0	0	0	0	0.3	0	0.8
TO4/6 (Cuorgnè)	0	0	0	0	0	0	0
TO5/1(Carmagnola)	0	0	0	0	0	0	0
TO5/2 (Chieri)	0	0	0	0	0.3	0	1
TO5/3 (Moncalieri)	0	0	0	0	0	0	0
TO5/4 (Nichelino)	0.4	0	1.3	1	0	0	0
VC/1(Vercelli-Santhià)	0.3	0	1	1	0	0	0
VC/2 (Montano-Borgosesia-Varallo)	0	0	0	0	0	0	0
VCO/1(Domodossola)	0	0	0	0	0	0	0
VCO/2 (Omegna)	0	0	0	0	0	0	0
VCO/3 (Verbania)	0.5	0	1.5	1	0.5	0	1.6

SURVIVAL

Median survival time and survival probability by calendar period

period	median (months)	SURVIVAL 1990-2017				
		6 months	1 yr	2 yrs	3 yrs	5 yrs
1990-1994	4.4	46%	23%	10%	8%	8%
1995-1999	4.7	36%	21%	16%	12%	5%
2000-2004	4.8	42%	23%	14%	7%	4%
2005-2009	6.1	51%	30%	17%	11%	9%
2010-2014	5.0	47%	32%	17%	15%	10%
2015-2017	2.4	34%	21%	17%	14%	-

Median survival time and survival probability by age class

age class	median (months)	SURVIVAL 1990-2016				
		6 months	1 yr	2 yrs	3 yrs	5 yrs
00-54	6.2	51%	37%	25%	22%	15%
55-64	8.0	54%	33%	22%	15%	10%
65-74	5.3	46%	28%	11%	7%	-
75+	2.0	23%	6%	1%	-	-

Median survival time and survival probability by histotype

histotype		SURVIVAL 1990-2017					
		median (months)	6 months	1 yr	2 yrs	3 yrs	5 yrs
epiteliomorfo		5.3	46%	27%	16%	11%	8%
bifasico		3.0	33%	15%	8%	-	-
sarcomatoso		2.8	18%	9%	9%	9%	-
non specificato		5.0	47%	30%	19%	16%	10%
non noto		1.7	14%	-	-	-	-